

Soft Γ - Modules

Serkan Onar¹, Sanem Yavuz², Bayram Ali Ersoy³

¹Yildiz Technical University, Department of Mathematics, Istanbul, Turkey
serkan10ar@gmail.com

²Yildiz Technical University, Department of Mathematics, Istanbul, Turkey
ssanemy@gmail.com

³Yildiz Technical University, Department of Mathematics, Istanbul, Turkey
ersoya@gmail.com

Received on: 14-04-2017. **Accepted** on: 28-06-2017. **Published** on: 30-06-2017

doi:10.23755/rm.v32i0.325

©Serkan Onar et al.



Abstract

In this paper, the definitions of soft Γ -module, soft Γ -module homomorphism and soft Γ -exactness are introduced with the aid of the concept of soft set theory introduced by Molodtsov. In the meantime, some of their properties and structural characteristics are investigated and discussed. Thereafter, several illustrative examples are given.

Keywords: soft set; soft module; Γ -ring; Γ -module; soft Γ -module; soft Γ -module homomorphism; soft Γ -module isomorphism; soft Γ -exactness.

2010 AMS subject classifications: 03E72; 08A72.

1 Introduction

In the real world, there are some various uncertainties but classical mathematical tools is not convenient for modeling these. Uncertain and unclear data which are contained by economy, engineering, environmental science, social science, medical science, business administration and many other fields are common. Although many diverse theories such as probability theory, soft set theory, intuitionistic fuzzy soft set theory and rough set theory are known and these present advantageous mathematical approaches for modeling of uncertainties, each of these theories have their inherent difficulties.

In 1999, Molodstov [1] developed soft set theory which is considered a mathematical tool for working with uncertainties. Since the emergence of soft set theory attracts attention and especially recently works on the soft set theory is progressing rapidly. Maji et al. [2] described some operations on soft sets and these operations are used soft sets of decision making problems. Chen et al. [3] offered a new definition for decrease of parametrization on soft sets. They made comparison between this definition and concept of restriction of property in the rough set theory. In theory, Maji et al. [4] worked various operator on soft set. Kong et al. [5] developed definition of parametrization reduction on soft set. Zou and Xiao suggested some approach of data analysis in case of insufficient information on soft set. Jiang et al. presented a unique approach of the semantic decision making by means of ontological thinking and ontology-based soft sets.

Besides studies on classic module theory have continued and interesting results have been discovered recently. Macias Diaz et al. [6] studied on modules which are isomorphic to relatively divisible or pure submodules of each other. Abuhlail et al. [7] presented on topological lattices and their applications to module theory. On the other hand, Ameri et al. [8] investigated gamma module and Davvaz et al. [9] studied tensor product of gamma modules.

As for soft module theory, Sun et al. [10] presented the notion of soft set and soft module. Xiang [11] worked soft module theory. T.Shah et al. [12] defined the notion of primary decomposition in a soft ring and soft module, and derived some related properties. Erami et al. [13] gave the concept of a soft MV-module and soft MV- submodule. In these days, there are some studies related with soft sets. Ali et al. [14] investigated some new operations in soft set theory and Pei et al. [15] studied from soft sets to information systems. Xiao et al. [16] presented research on synthetically evaluating method for business competitive capacity based on soft set. Aktaş et al. [17] showed soft sets and soft groups and Acar et al. [18] also showed soft sets and soft rings.

The main purpose of this paper is to deal with algebraic structure of Γ - module by applying soft set theory. The concept of soft Γ - module is introduced, their characterization and algebraic properties are investigated by giving some several

examples. In addition to this, soft Γ – homomorphism , soft Γ – isomorphism and their properties are introduced. After all, we make inferences that images of soft Γ – homomorphisms and inverse images of soft Γ – homomorphisms are soft Γ – homomorphisms. Furthermore soft Γ – exactness is investigated and illustrated with a related example.

2 Preliminaries

In this section, preliminary informations will be required to soft Γ – modules. First of all we give basic concepts of soft set theory.

Definition 2.1. [18] Let X denotes an initial universe set and E is a set of parameters. The power set of X is denoted by $P(X)$. A pair of (F, E) is called a soft set over X if and only if F is a mapping from E into the set of all subsets of X , i.e, $F: E \rightarrow P(X)$.

Definition 2.2. [18] Let (F, A) and (G, B) be two soft sets over a common universe X .

i) If $A \subseteq B$ and $F(a) \subseteq G(a)$ for all $a \in A$ then we say that (F, A) is a soft subset of (G, B) , denoted by $(F, A) \subseteq (G, B)$.

ii) If (F, A) is a soft subset of (G, B) and (G, B) is a soft subset of (F, A) , then we say that (F, A) is a soft equal to (G, B) , denoted by $(F, A) \cong (G, B)$.

Example 2.1. Let $X = M_2(Z_3)$ denotes an initial universe set, i.e, 2×2 matrices with Z_3 terms and $E = \left\{ \begin{bmatrix} \bar{0} & \bar{0} \\ \bar{0} & \bar{0} \end{bmatrix}, \begin{bmatrix} \bar{1} & \bar{0} \\ \bar{0} & \bar{1} \end{bmatrix} \right\}$ is a set of parameters. Then $F: E \rightarrow P(X)$ where $F\left(\begin{bmatrix} \bar{0} & \bar{0} \\ \bar{0} & \bar{0} \end{bmatrix}\right) = \left\{ \begin{bmatrix} \bar{0} & \bar{1} \\ \bar{1} & \bar{1} \end{bmatrix}, \begin{bmatrix} \bar{2} & \bar{1} \\ \bar{0} & \bar{2} \end{bmatrix} \right\}$, $F\left(\begin{bmatrix} \bar{1} & \bar{0} \\ \bar{0} & \bar{1} \end{bmatrix}\right) = \left\{ \begin{bmatrix} \bar{2} & \bar{0} \\ \bar{0} & \bar{2} \end{bmatrix} \right\}$. Clearly, (F, E) is called a soft set over X .

Definition 2.3. [18] Let (F, A) and (G, B) be two soft sets over a common universe X . The intersection of (F, A) and (G, B) is defined as the soft set (H, C) satisfying the following conditions:

i) $C = A \cap B$.

ii) For all $c \in C$, $H(c) = F(c)$ or $G(c)$.

In this case, we write $(F, A) \tilde{\cap} (G, B) = (H, C)$.

Definition 2.4. [18] Let (F, A) and (G, B) be two soft sets over a common universe X . The union of (F, A) and (G, B) is defined as the soft set (H, C) satisfying the following conditions:

i) $C = A \cup B$.

ii) For all $c \in C$,

$$H(c) = \left\{ \begin{array}{ll} F(c) & \text{if } c \in A - B, \\ G(c) & \text{if } c \in B - A, \\ F(c) \cup G(c) & \text{if } c \in A \cap B. \end{array} \right\}$$

This is denoted by $(F, A) \tilde{\cup} (G, B) = (H, C)$.

Definition 2.5. [18] If (F, A) and (G, B) are two soft sets over a common universe X , then (F, A) AND (G, B) denoted by $(F, A) \tilde{\wedge} (G, B)$ is defined as $(F, A) \tilde{\wedge} (G, B) = (H, C)$, where $C = A \times B$ and $H(x, y) = F(x) \cap G(y)$, for all $(x, y) \in C$.

Definition 2.6. Let $\{(F_i, A_i) : i \in I\}$ be a non- empty family soft sets. The \wedge -intersection of a non-empty family soft sets is defined by $(\psi, Y) = \tilde{\wedge}_{i \in I} (F_i, A_i)$ where (ψ, Y) is a soft set, $Y = \prod_{i \in I} A_i$ and $\psi(y) = \cap_{i \in I} F_i(y)$ for every $y = (y_i)_{i \in I} \in Y$.

Definition 2.7. [18] If (F, A) and (G, B) are two soft sets over a common universe X , then (F, A) OR (G, B) denoted by $(F, A) \tilde{\vee} (G, B)$ is defined as $(F, A) \tilde{\vee} (G, B) = (H, C)$, where $C = A \times B$ and $H(x, y) = F(x) \cup G(y)$, for all $(x, y) \in C$.

Definition 2.8. Let $\{(F_i, A_i) : i \in I\}$ be a non- empty family soft sets. The \vee -union of a non-empty family soft sets is defined by $(\psi, Y) = \tilde{\vee}_{i \in I} (F_i, A_i)$ where (ψ, Y) is a soft set, $Y = \prod_{i \in I} A_i$ and $\psi(y) = \cup_{i \in I} F_i(y)$ for every $y = (y_i)_{i \in I} \in Y$.

On the other hand we will introduce modules and soft modules, then we will study some properties and theories of soft modules such as trivial soft module, whole soft module, the concepts of soft submodule and soft module homomorphisms.

Definition 2.9. [10] Let R be a ring with identity. M is said to be a left R -module if left scalar multiplication $\lambda : R \times M \rightarrow M$ via $(a, x) \mapsto ax$ satisfying the axioms $\forall r, r_1, r_2, 1 \in R; m, m_1, m_2 \in M$:

- i) M is an abelian group,
- ii) $r(m_1 + m_2) = rm_1 + rm_2, (r_1 + r_2)m = r_1m + r_2m,$
- iii) $(r_1r_2)m = r_1(r_2m),$
- iv) $1m = m.$

Left R -module is denoted by ${}_R M$ or M for short. Similarly we can define right R - module and denote it by M_R .

Soft Γ - Modules

Example 2.2. Let $R = M_2(Z)$ and $M = \left\{ \begin{bmatrix} a \\ b \end{bmatrix} \mid a, b \in Z \right\}$. Then M is module on R .

Definition 2.10. [10] Let M be a left R - module, A be a any nonempty set and (F, A) is a soft set over M . (F, A) is said to be a soft module over M if and only if $F(x)$ is submodule over M , for all $x \in A$.

Definition 2.11. [10] Let (F, A) be a soft module over M then

i) (F, A) is said to be a trivial soft module over M if $F(x) = 0$ for all $x \in A$, where 0 is zero element of M .

ii) (F, A) is said to be an whole soft module over M if $F(x) = M$ for all $x \in A$.

Proposition 2.1. [10] Let (F, A) and (G, B) be two soft modules over M .

1) $(F, A) \widetilde{\cap} (G, B)$ is a soft module over M .

2) $(F, A) \widetilde{\cup} (G, B)$ is a soft module over M if $A \cap B = \emptyset$.

Definition 2.12. [10] If (F, A) and (G, B) be two soft modules over M , then $(F, A) + (G, B)$ is defined as $(H, A \times B)$, where $H(x, y) = F(x) + G(y)$ for all $(x, y) \in A \times B$.

Proposition 2.2. [10] Assume that (F, A) and (G, B) are two soft modules over M . Then $(F, A) + (G, B)$ is soft module over M .

Definition 2.13. [10] Suppose that (F, A) and (G, B) be two soft modules over M and N respectively. Then $(F, A) \times (G, B) = (H, A \times B)$ is defined as $H(x, y) = F(x) \times G(y)$ for all $(x, y) \in A \times B$.

Proposition 2.3. [10] Let (F, A) and (G, B) be two soft modules over M and N respectively. Then $(F, A) \times (G, B)$ is soft module over $M \times N$.

Definition 2.14. [10] Let (F, A) and (G, B) be two soft modules over M . Then (G, B) is soft submodule of (F, A) if

i) $B \subset A$,

ii) $G(x) < F(x), \forall x \in B$.

This is denoted by $(G, B) \widetilde{<} (F, A)$.

Proposition 2.4. [10] Let (F, A) and (G, B) be two soft modules over M . We say that (G, B) is soft submodule of (F, A) if $G(x) \subseteq F(x), \forall x \in A$.

Definition 2.15. [10] Assume that $E = \{e\}$, where e is unit of A . Then every soft module (F, A) over M at least have two soft modules (F, A) and (F, E) called trivial soft submodule.

Proposition 2.5. [10] Let (F, A) and (G, B) are two soft modules over M and (G, B) is soft submodule of (F, A) . If $f : M \rightarrow N$ is a homomorphism of module, then $(f(F), A)$ and $(f(G), B)$ are all soft modules over N and $(f(G), B)$ is soft submodule of $(f(F), A)$.

Definition 2.16. [10] Let (F, A) and (G, B) be two soft modules over M and N respectively, $f : M \rightarrow N, g : A \rightarrow B$ be two functions. Then we say that (f, g) is a soft homomorphism if the following conditions are satisfied:

i) $f : M \rightarrow N$ is a homomorphism of module,

ii) $g : A \rightarrow B$ is a mapping,

iii) For all $x \in A, f(F(x)) = G(g(x))$.

We say that (F, A) is a soft homomorphic to (G, B) which denoted by $(F, A) \tilde{\sim} (G, B)$. In this definition, if f is an isomorphism from M to N and g is a one-to-one mapping from A onto B , then we say that (F, A) is a soft isomorphism and that (F, A) is a soft isomorphic to (G, B) , this is denoted by $(F, A) \cong (G, B)$.

Finally, we will define Γ - ring and Γ - module and their homomorphisms which are basic definitions for soft Γ - module.

Definition 2.17. [8] Let R and Γ be additive abelian groups. Then we say that R is a Γ - ring if there exists a mapping:

$$\cdot : R \times \Gamma \times R \rightarrow R$$

$$(r_1, \gamma, r_2) \rightarrow r_1 \gamma r_2$$

such that for every $a, b, c \in R$ and $\alpha, \beta \in \Gamma$, the following hold:

i) $(a + b)\alpha c = a\alpha c + b\alpha c,$

ii) $a(\alpha + \beta)c = a\alpha c + a\beta c,$

iii) $a\alpha(b + c) = a\alpha b + a\alpha c,$

iv) $(a\alpha b)\beta c = a\alpha(b\beta c).$

Definition 2.18. [8] A subset A of a Γ - ring R is said to be a right ideal of R if A is an additive subgroup of R and $A\Gamma R \subseteq A$, where $A\Gamma R = \{a\alpha c \mid a \in A, \alpha \in \Gamma, r \in R\}$.

A left ideal of R is defined in a similar way. If A is both right and left ideal, we say that A is an ideal of R .

Definition 2.19. [8] If R and S are Γ - rings, then a pair (θ, φ) of maps from R into S is called a homomorphism from R into S if

i) $\theta(x + y) = \theta(x) + \theta(y),$

ii) φ is an isomorphism on $\Gamma,$

iii) $\theta(x\gamma y) = \theta(x)\varphi(\gamma)\theta(y).$

Definition 2.20. [8] Let R be a Γ - ring. A left Γ - module M is an additive abelian group M together with a mapping $\cdot : R \times \Gamma \times M \rightarrow M$ such that for all $m, m_1, m_2 \in M$ and $\gamma, \gamma_1, \gamma_2 \in \Gamma, r, r_1, r_2 \in R$ the following hold:

Soft Γ - Modules

- i)** $r\gamma(m_1 + m_2) = r\gamma m_1 + r\gamma m_2,$
- ii)** $(r_1 + r_2)\gamma m = r_1\gamma m + r_2\gamma m,$
- iii)** $r(\gamma_1 + \gamma_2)m = r\gamma_1 m + r\gamma_2 m,$
- iv)** $r_1\gamma_1(r_2\gamma_2 m) = (r_1\gamma_1 r_2)\gamma_2 m.$

A right Γ - module R is defined in analogous manner.

Example 2.3. Let $R = \{[\bar{k} \ \bar{m}] \mid k, m \in Z_2\}$, i.e, 1×2 matrices and $\Gamma = \left\{ \begin{bmatrix} \bar{0} \\ \bar{0} \end{bmatrix}, \begin{bmatrix} \bar{1} \\ \bar{0} \end{bmatrix} \right\} \in Z_2$, where Γ is 2×1 matrices. Then we say that R is a Γ - ring. Similarly, R and Γ are same if we choose $M = \{[\bar{0} \ \bar{0}], [\bar{1} \ \bar{1}]\}$, then M is Γ - module R .

Definition 2.21. [8] Presume that $(M, +)$ be an Γ - module R . A nonempty subset N of $(M, +)$ is said to be a left Γ - submodule R of M if N is a subgroup of M and $R\Gamma N \subseteq N$, where $R\Gamma N = \{r\gamma n \mid \gamma \in \Gamma, r \in R, n \in N\}$, that is for all $n_1, n_2 \in N$ and for all $\gamma \in \Gamma, r \in R; n_1 - n_2 \in N$ and $r\gamma n \in N$. In this case we write $N \leq M$.

Example 2.4. In previous example, let $N = \{[\bar{0} \ \bar{0}]\} \subset M$ and $H : N \rightarrow P(M)$ be a set valued function defined by $H(a) = \{b \in M \mid R(a, \alpha, b) \Leftrightarrow a\alpha b \in [\bar{0} \ \bar{0}]\}$ for all $a \in N$. H is clear that $H([\bar{0} \ \bar{0}]) = ([\bar{0} \ \bar{0}])$ is Γ - submodule R of M .

Definition 2.22. [8] Let M and N be arbitrary Γ - module R . A mapping $f : M \rightarrow N$ is a homomorphism of Γ - module R if for all $x, y \in M$ and $\forall r \in R, \forall \gamma \in \Gamma$ we have

- i)** $f(x + y) = f(x) + f(y),$
- ii)** $f(r\gamma x) = r\gamma f(x).$

A homomorphism f is monomorphism if f is one-to-one and f is epimorphism if f is onto. f is called isomorphism if f is both monomorphism and epimorphism. We denote the set of all R_Γ - homomorphisms from M into N by $Hom_{R_\Gamma}(M, N)$ or shortly by $Hom_{R_\Gamma}(M, N)$. In particular $M = N$ we denote $Hom(M, M)$ by $End(M)$.

Definition 2.23. [18] Let M be a nonempty set and a Γ -module. The pair (F, A) is a soft set over M . The set $Supp(F, A) = \{x \in A : F(x) \neq \emptyset\}$ is called a support of the soft set (F, A) . The soft set (F, A) is non-null if $Supp(F, A) \neq \emptyset$.

3 Soft Γ - Modules

In this section, firstly we will define soft Γ - modules, then we will give some operations on this modules. Throughout the section, M is a Γ -module.

Definition 3.1. Let (F, A) be a non-null soft set over M . Then, (F, A) is said to be a soft Γ -module over M if $F(a)$ is a Γ -submodule M such that $F : A \rightarrow P(M)$, (i.e. $a \rightarrow F(a)$) for all $a \in A, y \in \text{Supp}(F, A)$.

Example 3.1. For consider the additively abelian groups $\mathbb{Z}_6 = \{\bar{0}, \bar{1}, \bar{2}, \bar{3}, \bar{4}, \bar{5}\}$ and $\Gamma = \{\bar{0}, \bar{2}\}$. Let $\cdot : \mathbb{Z}_6 \times \Gamma \times \mathbb{Z}_6 \rightarrow \mathbb{Z}_6, (m_1, \Gamma, m_2) = m_1 \Gamma m_2$. Hence \mathbb{Z}_6 is a Γ -module. Let $A = \mathbb{Z}_6$ and $F : A \rightarrow P(M)$ be a set valued function defined by

$$\begin{aligned} f(\bar{0}) &= f(\bar{2}) = f(\bar{4}) = \mathbb{Z}_6, \\ f(\bar{1}) &= f(\bar{3}) = f(\bar{5}) = \{\bar{0}, \bar{3}\} \end{aligned}$$

are Γ -submodule of \mathbb{Z}_6 . Hence $(F, A,)$ is a soft Γ -module over \mathbb{Z}_6 .

Example 3.2. Let M is a Γ -module and (F, A) be a soft set over M . $F : A \rightarrow P(M)$ is defined by $F(x) = \{y \in M \mid x\alpha y = 0\}$ for all $x \in A, \alpha \in \Gamma$. It is clear that (F, A) is a soft Γ -module.

Example 3.3. For consider the additively abelian groups

$$\begin{aligned} M &= R = \{[\bar{0} \ \bar{0}], [\bar{1} \ \bar{0}], [\bar{0} \ \bar{1}], [\bar{1} \ \bar{1}]\} \subseteq (\mathbb{Z}_2)_{1 \times 2} \\ \text{and } \Gamma &= \left\{ \begin{bmatrix} \bar{0} \\ \bar{0} \end{bmatrix}, \begin{bmatrix} \bar{1} \\ \bar{0} \end{bmatrix} \right\} \subseteq (\mathbb{Z}_2)_{2 \times 1} \end{aligned}$$

with addition defined as matrix addition. It is trivial that R is a Γ -ring. Also M is a Γ -module over R . Let $N = \{[\bar{0} \ \bar{0}]\} \subseteq M$ and $H : N \rightarrow P(M)$ be a set valued function defined by $H(a) = \{b \in M \mid R(a, \alpha, b) \leftrightarrow a\alpha b \in [\bar{0} \ \bar{0}], \forall \alpha \in \Gamma\}$ for all $a \in N$. It is clear that $H([\bar{0} \ \bar{0}]) = \{[\bar{0} \ \bar{0}]\}$ are sub Γ -module of M . Hence (H, N) is soft Γ -module of M .

Theorem 3.1. Let (F, A) and (G, B) are two soft Γ -modules over M . Then $(F, A) \tilde{\cap} (G, B)$ is a soft Γ -module over M if it is non-null.

Proof. By definition, we have that $(F, A) \tilde{\cap} (G, B) = (H, C)$ where $H(c) = F(x) \cap G(y)$ for all $c \in C$. We assume that (H, C) is a non-null soft set over M . If $c \in \text{Supp}(H, C)$, then $H(c) = F(x) \cap G(y) \neq \emptyset$. We know that $(F, A), (G, B)$ are both soft Γ -module over M , and so, the nonempty sets $F(x)$ and $G(y)$ are both Γ -submodule over M . Thus, $H(c)$ is a Γ -submodule over M for all $c \in \text{Supp}(H, C)$. In this position, $(H, C) = (F, A) \tilde{\cap} (G, B)$ is a soft Γ -module over M . \square

Theorem 3.2. Let (F, A) and (G, B) are two soft Γ -modules over M . Then $(F, A) \tilde{\cup} (G, B)$ is a soft Γ -module over M if $A \cap B = \emptyset$.

Proof. By definition, we have that $(F, A) \tilde{\cup} (G, B) = (H, C)$ where $H(c) = F(x) \cap G(y)$ for all $c \in C$. Note first that (H, C) is a non-null owing to the fact that $Supp(H, C) = Supp(F, A) \tilde{\cup} (G, B)$. Suppose that $c \in Supp(H, C)$. Then $H(c) \neq \emptyset$ so we have $F(x), G(y) \neq \emptyset$. From the hypothesis $A \cap B = \emptyset$, we follow that $H(c) = F(x) \cap G(y)$. On the other hand $F(x) \cap G(y)$ is a soft Γ -module over M , we conclude that (H, C) is a soft Γ -module over M for all $c \in Supp(H, C)$. Consequently $(F, A) \tilde{\cup} (G, B) = (H, C)$ is a soft Γ -module over M . \square

On the other hand, union of two soft Γ -modules is not always soft Γ -module. We will explain this situation with following example.

Example 3.4. Let $M = \mathbb{Z}_6 = \{\bar{0}, \bar{1}, \bar{2}, \bar{3}, \bar{4}, \bar{5}\}$ is a M_Γ -module, $\Gamma = \{\bar{0}, \bar{1}\}$, $A = \mathbb{Z}_2 = \{\bar{0}, \bar{1}\}$ and $B = \mathbb{Z}_3 = \{\bar{0}, \bar{1}, \bar{2}\}$ such that $F(\bar{0}) = F(\bar{1}) = \{\bar{0}, \bar{2}, \bar{4}\}$, $G(\bar{0}) = G(\bar{1}) = G(\bar{2}) = \{\bar{0}, \bar{3}\}$, $A \cap B = \{\bar{0}, \bar{1}\}$. If this condition is hold, then $(F, A) \tilde{\cup} (G, B)$ is not a soft Γ -module over M . Indeed, $H(\bar{1}) = \{\bar{0}, \bar{2}, \bar{3}, \bar{4}\} \notin P(M)$.

Definition 3.2. If (F, A) and (G, B) are two soft Γ -modules over M , then (F, A) AND (G, B) denoted by $(F, A) \tilde{\wedge} (G, B)$ is defined as $(F, A) \tilde{\wedge} (G, B) = (H, C)$, where $C = A \times B$ and $H(x, y) = F(x) \tilde{\cap} G(y)$, for all $(x, y) \in C$.

Theorem 3.3. Suppose that (F, A) and (G, B) are two soft Γ -modules over M . Then $(F, A) \tilde{\wedge} (G, B)$ is soft Γ -module over M if it is non-null.

Proof. Using definition, we have that $(F, A) \tilde{\wedge} (G, B) = (H, C)$ where $C = A \times B$ and $H(x, y) = F(x) \tilde{\cap} G(y)$, for all $(x, y) \in C$. Then the hypothesis, (H, C) is a non-null soft set over M . Since (H, C) is a non-null, $Supp(H, C) \neq \emptyset$ and so, for $(x, y) \in Supp(H, C)$, $H(x, y) = F(x) \tilde{\cap} G(y) \neq \emptyset$. We assume that $t_1, t_2 \in F(x) \tilde{\cap} G(y)$. In this position

i) If $t_1, t_2 \in F(x) = \{y : R(x, y)\}$ we have that $xt_1 \in A, xt_2 \in A$. This implies that $x(t_1 + t_2) \in A$.

ii) If $t_1, t_2 \in G(y) = \{y_1 : R(y, y_1)\}$ we have that $yt_1 \in B, yt_2 \in B$. This implies that $y(t_1 + t_2) \in B$.

Hence $F(x) \tilde{\cap} G(y)$ is a Γ -submodule. By the definition of soft Γ -module, (F, A) and (G, B) are soft Γ -modules over M . $F(x), G(y)$ are also Γ -submodule over M . Furthermore $H(x, y) = F(x) \tilde{\cap} G(y)$ is a Γ -submodule over M for all $(x, y) \in (H, C) = (F, A) \tilde{\wedge} (G, B)$. Hence $(F, A) \tilde{\wedge} (G, B)$ is soft Γ -module over M . \square

Definition 3.3. If (F, A) and (G, B) are two soft Γ -modules over M , then (F, A) OR (G, B) denoted by $(F, A) \tilde{\vee} (G, B)$ is defined as $(F, A) \tilde{\vee} (G, B) = (H, C)$, where $C = A \times B$ and $H(x, y) = F(x) \tilde{\cup} G(y)$, for all $(x, y) \in C$.

Theorem 3.4. Suppose that (F, A) and (G, B) are two soft Γ -modules over M . Then $(F, A) \tilde{\vee} (G, B)$ is soft Γ -module over M .

Proof. Using definition, we have that $(F, A) \tilde{\vee} (G, B) = (H, C)$, where $C = A \times B$ and $H(x, y) = F(x) \tilde{\cup} G(y)$, for all $(x, y) \in C$. Assume that $c \in \text{Supp}(H, C)$. Then $H(c) \neq \emptyset$ and so we have that $F(x) \neq \emptyset, G(y) \neq \emptyset$. By assumption, $F(x) \tilde{\cup} G(y)$ is a soft Γ -module of M for all $c \in \text{Supp}(H, C)$. Consequently $(F, A) \tilde{\vee} (G, B) = (H, C)$ is a soft Γ -module over M . \square

Definition 3.4. Let (F, A) and (G, B) are two soft Γ -modules over M . Then $(F, A) \tilde{+} (G, B) = (H, A \times B)$ is defined as $H(x, y) = F(x) + G(y)$ for all $(x, y) \in A \times B$.

Theorem 3.5. Suppose that (F, A) and (G, B) are two soft Γ -modules over M . Then $(F, A) \tilde{+} (G, B)$ is soft Γ -module over M .

Proof. By the definition we write $(F, A) \tilde{+} (G, B) = (H, A \times B)$ and $H(x, y) = F(x) + G(y)$ for all $(x, y) \in A \times B$. Let $(x, y) \in \text{Supp}(H, A \times B)$. Then, $H(x, y) \neq \emptyset$ and so we have $F(x) \neq \emptyset, G(y) \neq \emptyset$. By taking into account, (F, A) and (G, B) are two soft Γ -modules over M , it follows that $F(x) + G(y)$ is a soft Γ -module over M for all $(x, y) \in \text{Supp}(H, A \times B)$. Hence $(F, A) \tilde{+} (G, B)$ is soft Γ -module over M . \square

Definition 3.5. Let (F, A) and (G, B) are two soft Γ -modules over M . Then $(F, A) \tilde{\times} (G, B) = (H, A \times B)$ is defined as $H(x, y) = F(x) \times G(y)$ for all $(x, y) \in A \times B$.

Theorem 3.6. Suppose that (F, A) and (G, B) are two soft Γ -modules over M . Then $(F, A) \tilde{\times} (G, B)$ is soft Γ -module over M .

Proof. By the definition we write $(F, A) \tilde{\times} (G, B) = (H, A \times B)$ and $H(x, y) = F(x) \times G(y)$ for all $(x, y) \in A \times B$. Let $(x, y) \in \text{Supp}(H, A \times B)$. Then, $H(x, y) \neq \emptyset$ and so we have $F(x) \neq \emptyset, G(y) \neq \emptyset$. By taking into account, (F, A) and (G, B) are two soft Γ -modules over M , it follows that $F(x) \times G(y)$ is a soft Γ -module over M for all $(x, y) \in \text{Supp}(H, A \times B)$. Hence $(F, A) \tilde{\times} (G, B)$ is soft Γ -module over M . \square

Definition 3.6. Let (F, A) and (G, B) are two soft Γ -modules over M . Then (G, B) is called a soft Γ -submodule of (F, A) if

- i) $B \subseteq A$,
- ii) $\forall b \in \text{Supp}(G, B), g(b)$ is a Γ -submodule of $F(b)$.

This denoted by $(G, B) \subset (F, A)$. From the definition, it is easily deduced that if (G, B) is a soft Γ -submodule of (F, A) , then $\text{Supp}(G, B) \subset \text{Supp}(F, A)$.

Theorem 3.7. Let (F, A) and (G, B) be two soft Γ -modules over M and $(F, A) \tilde{\subseteq} (G, B)$. Then $(G, B) \subset (F, A)$.

Proof. Straight forward. \square

Corolary 3.1. Let (F, A) be a soft Γ -module over M and $\{(F_i, A_i) : i \in I\}$ be a nonempty family of soft Γ -submodules of (F, A) . Then,

- i) $\widetilde{\cap}_{i \in I}(F_i, A_i)$ is a soft Γ -submodule of (F, A) if it is non-null.
- ii) $\widetilde{\cup}_{i \in I}(F_i, A_i)$ is a soft Γ -submodule of (F, A) , if $A_i \cap A_j = \emptyset$ for all $i, j \in I$ and if it is non-null.
- iii) If $F_i(a_i) \subseteq F_j(a_j)$ or $F_j(a_j) \subseteq F_i(a_i)$ for all $i, j \in I, a_i \in A_i$, then $\widetilde{\vee}_{i \in I}(F_i, A_i)$ is a soft Γ -submodule of $\widetilde{\vee}_{i \in I}(F, A)$.
- iv) $\widetilde{\wedge}_{i \in I}(F_i, A_i)$ is a soft Γ -submodule of $\widetilde{\wedge}_{i \in I}(F, A)$.
- v) The cartesian product of the family $\prod_{i \in I}(F_i, A_i)$ is a soft Γ -submodule of $\prod_{i \in I}(F, A)$.
- vi) $\widetilde{\sum}_{i \in I}(F_i, A_i)$ is a soft Γ -submodule of $\widetilde{\sum}_{i \in I}(F, A)$.

Proof. Similar to the proof of Theorems 3.5, 3.6, 3.9, 3.11, 3.13 and 3.15. \square

4 Soft Γ -Module Homomorphism

In this section, firstly we will define trivial and whole soft Γ -modules over Γ -module M , homomorphism of Γ -modules and their properties. Moreover we will study soft Γ -module homomorphism and soft Γ -module isomorphism. Throughout the section, M is a Γ -module.

Definition 4.1. Let (ρ, A) and (σ, B) be two soft Γ -modules over Γ -module M and Γ -module M_1 respectively. Let $f : M \rightarrow M_1$ and $g : A \rightarrow B$ be two functions. The following conditions:

- i) f is an epimorphism of Γ -module,
 - ii) g is a surjective mapping,
 - iii) $f(\rho(y)) = \sigma(\rho(y))$ for all $y \in A$,
- were satisfied by the pair (f, g) , then (f, g) is called soft Γ -module homomorphism.

If there exists a soft Γ -module homomorphism between (ρ, A) and (σ, B) , we say that (ρ, A) is soft homomorphic to (σ, B) , and is denoted by $(\rho, A) \sim (\sigma, B)$. If there exists a soft Γ -module isomorphism between (ρ, A) and (σ, B) , we say that (ρ, A) is soft isomorphic to (σ, B) , and is denoted by $(\rho, A) \widetilde{\sim} (\sigma, B)$.

Definition 4.2. Let (F, A) be soft Γ -module over M .

- i) (F, A) is called the trivial soft Γ -module over M if $F(a) = \{0\}$ for all $a \in A$.

ii) (F, A) is called the whole soft Γ -module over M if $F(a) = M$ for all $a \in A$.

Definition 4.3. Let M and M_1 be two Γ -modules and $m : M \rightarrow M_1$ a mapping of Γ -module. If (F, A) and (G, B) are soft sets over M and M_1 respectively, then

i) $(m(F), A)$ is a soft set over M_1 where $m(F) : A \rightarrow P(M_1)$, $m(F)(a) = m(F(a))$ for all $a \in A$.

ii) $(m^{-1}(G), B)$ is a soft set over M where $m^{-1}(G) : B \rightarrow P(M)$, $m^{-1}(G)(b) = m^{-1}(G(b))$ for all $b \in B$.

Corolary 4.1. Let $m : M \rightarrow M_1$ be an onto homomorphism of Γ -module. Then following statements can be given.

i) (F, A) be soft Γ -module over M , then $(m(F), A)$ is a soft Γ -module over Γ -module M_1 .

ii) (G, B) be soft Γ -module over Γ -module M_1 , then $(m^{-1}(G), B)$ is a soft Γ -module over M .

Proof. i) Since (F, A) is a soft Γ -module over M , it is clear that $(m(F), A)$ is a non-null soft set over M_1 . For every $y \in \text{Supp}(m(F), A)$ we have $m(F)(y) = m(F(y)) \neq \emptyset$. Hence $m(F(y))$ which is the onto homomorphic image of Γ -module $F(y)$ is a Γ -module of M_1 for all $y \in \text{Supp}(m(F), A)$. That is $(m(F), A)$ is a soft Γ -module over Γ -module M_1 .

ii) It is easy to see that $\text{Supp}(m^{-1}(G), B) \subseteq \text{Supp}(G, B)$. By this way let $y \in \text{Supp}(m^{-1}(G), B)$. Then, $G(y) \neq \emptyset$. Hence $m^{-1}(G(y))$ which is homomorphic inverse image of Γ -module $G(y)$, is a soft Γ -module over M for all $y \in B$. \square

Theorem 4.1. Let $m : M \rightarrow M_1$ be a homomorphism of Γ -module and (F, A) , (G, B) be two soft Γ -modules over Γ -module M and Γ -module M_1 respectively. Then following statements can be given.

i) If $F(a) = \ker(m)$ for all $a \in A$, then $(m(F), A)$ is the trivial soft Γ -module over M_1 .

ii) If m is onto and (F, A) is whole, then $(m(F), A)$ is the whole soft Γ -module over M_1 .

iii) If $G(b) = m(M)$ for all $b \in B$, then $(m^{-1}(G), B)$ is the whole soft Γ -module over M .

iv) If m is injective and (G, B) is trivial, then $(m^{-1}(G), B)$ is the trivial soft Γ -module over M .

Proof. i) By using $F(a) = \ker(m)$ for all $a \in A$. Then $m(F)(a) = m(F(a)) = \{0_{M_1}\}$ for all $a \in A$. Hence $(m(F), A)$ is soft Γ -module over M_1 .

Soft Γ - Modules

ii) Suppose that m is onto and (F, A) is whole. Then $F(a) = M$ for all $a \in A$ and so $m(F)(a) = m(F(a)) = m(M) = M_1$ for all $a \in A$. Hence $(m(F), A)$ is whole soft Γ -module over M_1 .

iii) If we use hypothesis $G(b) = m(M)$ for all $b \in B$, we can write $m^{-1}(G)(b) = m^{-1}(G(b)) = m^{-1}(m(M)) = M$ for all $b \in B$. It is clear that, $(m^{-1}(G), B)$ is the whole soft Γ -module over M .

iv) Suppose that m is injective and (G, B) is trivial. Then, $G(b) = \{0\}$ for all $b \in B$, so $m^{-1}(G)(b) = m^{-1}(G(b)) = m^{-1}(\{0\}) = \ker m = \{0_M\}$ for all $b \in B$. Consequently, $(m^{-1}(G), B)$ is the trivial soft Γ -module over M . \square

Theorem 4.2. *Let $m : M \rightarrow M_1$ be a homomorphism of Γ -module and (F, A) , (G, B) be two soft Γ -modules over M . If (G, B) is soft Γ -submodule of (F, A) , then $(m(G), B)$ is soft Γ -submodule of $(m(F), A)$.*

Proof. Suppose that $y \in \text{Supp}(G, B)$. Then $y \in \text{Supp}(F, A)$. We know that $B \subseteq A$ and $G(y)$ is a Γ -submodule $F(y)$ for all $y \in \text{Supp}(G, B)$. From the expression hypothesis m is a homomorphism, $m(G)(y) = m(G(y))$ is a Γ -submodule of $m(F)(y) = m(F(y))$ and therefore $(m(G), B)$ is soft Γ -submodule of $(m(F), A)$. \square

Theorem 4.3. *Let $m : M \rightarrow M_1$ be a homomorphism of Γ -module and (F, A) , (G, B) be two soft Γ -modules over M . If (G, B) is soft Γ -submodule of (F, A) , then $(m^{-1}(G), B)$ is soft Γ -submodule of $(m^{-1}(F), A)$.*

Proof. Let $y \in \text{Supp}(m^{-1}(G), B)$. $B \subseteq A$ and $G(y)$ is a Γ -submodule of $F(y)$ for all $y \in B$. Since m is a homomorphism, $m^{-1}(G)(y) = m^{-1}(G(y))$ is a Γ -submodule of $m^{-1}(G(y)) = m^{-1}(G(y))$ for all $y \in \text{Supp}(m^{-1}(G), B)$. Hence $(m^{-1}(G), B)$ is soft Γ -submodule of $(m^{-1}(F), A)$. \square

5 Soft Γ - Exactness

In this section, we will introduce maximal and minimal soft Γ -submodules. Then, we will investigate short exact and exact sequence of Γ -modules. Finally, we will explain soft Γ -exactness and some their basic theories. Throughout this section M is Γ -module.

Definition 5.1. *Let (F, A) and (G, B) be two soft Γ -modules over M and (G, B) be soft Γ -submodule of (F, A) . We say (G, B) is maximal soft Γ -submodule of (F, A) if $G(x)$ is a maximal Γ -submodule of $F(x)$ for all $x \in B$. We say (G, B) is minimal soft Γ -submodule of (F, A) if $G(x)$ is a minimal Γ -submodule of $F(x)$ for all $x \in B$.*

Proposition 5.1. *Let (F, A) be a soft Γ -module over M .*

i) If $\{(G_i, B_i) \mid i \in I\}$ is a nonempty family of maximal soft Γ -submodules of (F, A) , then $\bigcap_{i \in I} (G_i, B_i)$ is maximal soft Γ -submodule of (F, A) .

ii) If $\{(G_i, B_i) \mid i \in I\}$ is a nonempty family of minimal soft Γ -submodules of (F, A) , then $\sum_{i \in I} (G_i, B_i)$ is minimal soft Γ -submodule of (F, A) .

Proof. straight forward. \square

Corolary 5.1. *Let (F, A) be a soft Γ -module over M and $f : M \rightarrow N$ be a homomorphism if $F(x) = \ker f$ for all $x \in A$, then $(f(F), A)$ is the rival soft Γ -module over N . Similarly, let (F, A) be an whole soft Γ -module over M and $f : M \rightarrow N$ be an epimorphism, then $(f(F), A)$ is a whole soft Γ -module over N .*

Definition 5.2. *The homomorphism sequence of Γ -modules $\dots \rightarrow M_{n-1} \xrightarrow{f_{n-1}} M_n \xrightarrow{f_n} M_{n+1} \rightarrow \dots$ is called exact sequence of Γ -modules if $\text{Im} f_{n-1} = \text{Ker} f_n$ for all $n \in \mathbb{N}$ and we call the exact sequence of Γ -modules form as $0 \rightarrow M_1 \xrightarrow{f} M \xrightarrow{g} M_2 \rightarrow 0$ the short exact sequence of Γ -modules.*

Proposition 5.2. *Let (F, A) be a trivial soft Γ -module over Γ -module M_1 and (G, B) be a whole soft Γ -module over Γ -module M_2 if $0 \rightarrow M_1 \xrightarrow{f} M \xrightarrow{g} M_2 \rightarrow 0$ is a short exact sequence, then $0 \rightarrow F(x) \xrightarrow{\tilde{f}} M \xrightarrow{\tilde{g}} G(y) \rightarrow 0$ is a short exact sequence for all $x \in A, y \in B$.*

Proof. $F(x) = 0, \forall x \in A$ since (F, A) is a trivial soft Γ -module over Γ -module M_1 , so \tilde{f} is a monomorphism. $G(y) = M_2, \forall y \in B$ since (G, B) is a whole soft Γ -module over Γ -module M_2 . $g : M \rightarrow M_2$ is an epimorphism as $0 \rightarrow M_1 \xrightarrow{f} M \xrightarrow{g} M_2 \rightarrow 0$ is a short exact sequence, so \tilde{g} is an epimorphism. \square

Proposition 5.3. *Let (F, A) be a trivial soft Γ -module over Γ -module M_1 and (G, B) be a whole soft Γ -module over Γ -module M if $0 \rightarrow M_1 \xrightarrow{f} M \xrightarrow{g} M_2 \rightarrow 0$ is a short exact sequence, then $0 \rightarrow f(F)(x) \xrightarrow{\tilde{f}} M \xrightarrow{\tilde{g}} g(G)(y) \rightarrow 0$ is a short exact sequence for all $x \in A, y \in B$.*

Proof. $F(x) = 0, \forall x \in A$ since (F, A) is a trivial soft Γ -module over Γ -module M_1 . $\text{Ker} f = 0$, so $\text{Ker} f = F(x), \forall x \in A$, consequently $(f(F), A)$ is trivial soft Γ -module over M . (G, B) is a whole soft Γ -module over M and $g : M \rightarrow M_2$ is an epimorphism, so $(g(G), B)$ is a whole soft Γ -module over M_2 , thus $0 \rightarrow f(F)(x) \xrightarrow{\tilde{f}} M \xrightarrow{\tilde{g}} g(G)(y) \rightarrow 0$ is a short exact sequence for all $x \in A, y \in B$. \square

Soft Γ - Modules

Definition 5.3. Let (F, A) , (G, B) and (H, C) are three soft Γ -modules over Γ -modules M, N and K respectively. Then we say soft Γ - exactness at (G, B) , if the following conditions are satisfied:

- i) $M \xrightarrow{f_1} N \xrightarrow{f_2} K$ is exact,
 - ii) $A \xrightarrow{g_1} B \xrightarrow{g_2} C$ is exact,
 - iii) $f_1(F(x)) = G(g_1(x))$ for all $x \in A$,
 - iv) $f_2(G(x)) = H(g_2(x))$ for all $x \in B$,
- which is denoted by $(F, A) \xrightarrow{(f_1, g_1)} (G, B) \xrightarrow{(f_2, g_2)} (H, C)$.

In this definition, if every $(F_i, A_i), i \in I$ is soft Γ - exact, then we say that $(F_i, A_i)_{i \in I}$ is soft Γ - exact.

Proposition 5.4. Let (F, A) and (G, B) are two soft Γ -modules over Γ -modules M and N respectively. If $(F, A) \xrightarrow{(f, g)} (G, B) \rightarrow 0$ is soft Γ - exact, then (f, g) is soft Γ - homomorphism. In particular, if $0 \rightarrow (F, A) \xrightarrow{(f, g)} (G, B) \rightarrow 0$ is soft Γ - exact, then (f, g) is soft Γ -isomorphism.

Proof. Since $(F, A) \xrightarrow{(f, g)} (G, B) \rightarrow 0$ is soft Γ - exact, we have $M \xrightarrow{f} N \rightarrow 0$ and $A \xrightarrow{g} B \rightarrow 0$ are exact. Thus f and g are epimorphisms, it is clear that (f, g) is homomorphism. If $0 \rightarrow (F, A) \xrightarrow{(f, g)} (G, B) \rightarrow 0$ is soft Γ - exact, then $0 \rightarrow M \xrightarrow{f} N \rightarrow 0$ and $0 \rightarrow A \xrightarrow{g} B \rightarrow 0$ are exact. Thus f and g are isomorphisms, it is clear that (f, g) is soft Γ -isomorphism. \square

Definition 5.4. Let $M = 0$ and $A = 0$, then $(F, A) = 0$. We call (F, A) is a zero-soft Γ - module.

Proposition 5.5. Let (F, A) , (G, B) and (H, C) are three soft Γ -modules over Γ -modules M, N and K respectively. If $(F, A) \xrightarrow{(f_1, g_1)} (G, B) \xrightarrow{(f_2, g_2)} (H, C)$ is soft Γ - exact with f_1, g_1 epimorphism and f_2, g_2 monomorphism, then (G, B) is a zero-soft Γ - module.

Proof. Since $(F, A) \xrightarrow{(f_1, g_1)} (G, B) \xrightarrow{(f_2, g_2)} (H, C)$ is soft Γ - exact with f_1, g_1 epimorphism and f_2, g_2 monomorphism, we have $M \xrightarrow{f_1} N \xrightarrow{f_2} K$ and $A \xrightarrow{g_1} B \xrightarrow{g_2} C$, hence $N = 0$ and $B = 0$, it is clear that (G, B) is zero-soft Γ - module. \square

Theorem 5.1. Let (F, A) and (H, B) are two soft Γ -modules over Γ -modules M and N respectively. For any $M \subset N, A \subset B$ and $M \subset H(x)$ where $x \in B$. If $(F, A) \xrightarrow{(f, g)} (H, B)$ is soft Γ -homomorphism, then $0 \rightarrow (F, A) \xrightarrow{(f, g)} (H, B) \xrightarrow{(f_1, g_1)} (I, B/A) \rightarrow 0$ is soft Γ - exact, where $I(x + A) = H(x)/M$ for all $x \in B$.

Proof. We know that $0 \rightarrow M \xrightarrow{f} N \xrightarrow{f_1} N/M \rightarrow 0$ and $0 \rightarrow A \xrightarrow{g} B \xrightarrow{g_1} B/A \rightarrow 0$ are exact. It is clear that M is a Γ -submodule of N , so that N/M is a Γ -module and M is a Γ -submodule of $H(x)$ and $H(x)/M$ is always a Γ -submodule of N/M . This shows that $(I, B/A)$ is a soft Γ -module over N/M . For all $x \in B/A$. Define $f_1 : N \rightarrow N/M$ by $f_1(n) = n + M$, for all $n \in N$. Meanwhile, we define $g_1 : B \rightarrow B/A$ by $g_1(b) = b + A$, for all $b \in B$. Therefore, it gives that

$$f_1(H(x)) = H(x) + M, I(g_1(x)) = I(x + A) = H(x) + M$$

for all $x \in B$, and hence $f_1(H(x)) = I(g_1(x))$. This implies

$$0 \rightarrow (F, A) \xrightarrow{(f,g)} (H, B) \xrightarrow{(f_1,g_1)} (I, B/A) \rightarrow 0$$

is soft Γ -exact. \square

Theorem 5.2. Let $(F, A_2), (G, A_1)$ and (H, A) are three soft Γ -modules over Γ -modules M_2, M_1 and M respectively. If M_1 and M_2 are Γ -submodules of M with $M_2 \subset M_1, A_1$ and A_2 are Γ -submodules of A with $A_2 \subset A_1$, where $M_1 \subset H(x)$, for all $x \in A$ and $M_2 \subset G(x)$ for all $x \in A_1$. Then $0 \rightarrow (I, A_1/A_2) \xrightarrow{(f_1,g_1)} (J, A/A_1) \xrightarrow{(f_2,g_2)} (P, A/A_1) \rightarrow 0$ is soft Γ -exact, where $I(x + A_2) = G(x)/M_2$, for all $x \in A_1$, $J(x + A_2) = H(x)/M_2$, for all $x \in A$, $P(x + A_1) = H(x)/M_1$, for all $x \in A$.

Proof. Since M_1 and M_2 are Γ -submodules of M with $M_2 \subset M_1$, we have a short exact sequence $0 \rightarrow M_1/M_2 \xrightarrow{f_1} M/M_2 \xrightarrow{f_2} M/M_1 \rightarrow 0$. Since A_1 and A_2 are Γ -submodules of A with $A_2 \subset A_1$, there is a short exact sequence $0 \rightarrow A_1/A_2 \xrightarrow{g_1} A/A_2 \xrightarrow{g_2} A/A_1 \rightarrow 0$. It is clear that M_2 is a Γ -submodule of M_1 , so that M_1/M_2 is a Γ -module. It gives that $G(x)/M_2$ is a Γ -module for all $x \in A_1$ from M_2 is a Γ -submodule of $G(x)$. However $G(x)/M_2$ is always a Γ -submodule of M_1/M_2 . This shows that $(I, A_1/A_2)$ is a soft Γ -module over M_1/M_2 for all $x \in A_1/A_2$. It is clear that $(J, A/A_2)$ and $(P, A/A_1)$ be a soft Γ -module over M/M_2 and M/M_1 respectively.

Define $f_1 : M_1/M_2 \rightarrow M/M_2$ by $f_1(m_1 + M_2) = m + M_2$, for all $m_1 \in M_1$. Meanwhile, we define $g_1 : A_1/A_2 \rightarrow A/A_2$ by $g_1(a_1 + A_2) = a + A_2$, for all $a_1 \in A_1$. Therefore, we have $f_1(I(x)) = f_1(G(x)/M_2) = H(x) + M_2, J(g_1(x)) = J(x + A_2) = H(x) + M_2$ for all $x \in A_1/A_2$, so $f_1(I(x)) = J(g_1(x))$ for all $x \in A_1/A_2$.

Define $f_2 : M/M_2 \rightarrow M/M_1$ by $f_2(m + M_2) = m + M_1$, for all $m \in M$. Let $g_2 : A/A_2 \rightarrow A/A_1$ be defined by $g_2(a + A_2) = a + A_1$, for all $a \in A$. Also, we have $f_2(J(x)) = f_2(H(x)/M_2) = H(x) + M_1$ for all $x \in A/A_2$, so $f_2(J(x)) = P(g_2(x))$ for all $x \in A/A_2$. Hence $0 \rightarrow (I, A_1/A_2) \xrightarrow{(f_1,g_1)} (J, A/A_1) \xrightarrow{(f_2,g_2)} (P, A/A_1) \rightarrow 0$ is soft Γ -exact. \square

Theorem 5.3. Let $(F_i, A_i), i = 1, 2, 3, 4, 5$ be a soft Γ -module over Γ -module $M_i, i = 1, 2, 3, 4, 5$ respectively. If $0 \rightarrow (F_1, A_1) \xrightarrow{(f_1, g_1)} (F_2, A_2) \xrightarrow{(f_2, g_2)} (F_3, A_3) \rightarrow 0$ and $0 \rightarrow (F_3, A_3) \xrightarrow{(f_3, g_3)} (F_4, A_4) \xrightarrow{(f_4, g_4)} (F_5, A_5) \rightarrow 0$ are soft Γ -exact. Then $0 \rightarrow (F_1, A_1) \xrightarrow{(f_1, g_1)} (F_2, A_2) \xrightarrow{(f_3 f_2, g_3 g_2)} (F_4, A_4) \xrightarrow{(f_4, g_4)} (F_5, A_5) \rightarrow 0$ is soft Γ -exact.

Proof. Since $0 \rightarrow (F_1, A_1) \xrightarrow{(f_1, g_1)} (F_2, A_2) \xrightarrow{(f_2, g_2)} (F_3, A_3) \rightarrow 0$ and $0 \rightarrow (F_3, A_3) \xrightarrow{(f_3, g_3)} (F_4, A_4) \xrightarrow{(f_4, g_4)} (F_5, A_5) \rightarrow 0$ are soft Γ -exact, we have $0 \rightarrow M_1 \xrightarrow{f_1} M_2 \xrightarrow{f_2} M_3 \rightarrow 0$ and $0 \rightarrow M_3 \xrightarrow{f_3} M_4 \xrightarrow{f_4} M_5 \rightarrow 0$ are exact. It is clear that $0 \rightarrow M_1 \xrightarrow{f_1} M_2 \xrightarrow{f_3 f_2} M_4 \xrightarrow{f_4} M_5 \rightarrow 0$ is exact. Since $0 \rightarrow A_1 \xrightarrow{g_1} A_2 \xrightarrow{g_2} A_3 \rightarrow 0$ and $0 \rightarrow A_3 \xrightarrow{g_3} A_4 \xrightarrow{g_4} A_5 \rightarrow 0$ are exact. It is clear that $0 \rightarrow A_1 \xrightarrow{g_1} A_2 \xrightarrow{g_3 g_2} A_4 \xrightarrow{g_4} A_5 \rightarrow 0$ is exact. Since $f_2(F_2(x)) = F_3(g_2(x))$ for all $x \in A_2$ and $f_3(F_3(x)) = F_4(g_3(x))$ for all $x \in A_3$. We have $f_3 f_2(F_2(x)) = f_3(F_3(g_2(x))) = F_4(g_3 g_2(x))$ for all $x \in A_2$. This implies $0 \rightarrow (F_1, A_1) \xrightarrow{(f_1, g_1)} (F_2, A_2) \xrightarrow{(f_3 f_2, g_3 g_2)} (F_4, A_4) \xrightarrow{(f_4, g_4)} (F_5, A_5) \rightarrow 0$ is soft Γ -exactness. \square

6 Conclusion

In this work the theoretical point of view of soft Γ - module is discussed. The work is focused on soft Γ - module, soft Γ - module homomorphism and soft Γ -exactness. By using these concepts, we studied the algebraic properties of soft sets in Γ - module structure. One could extend this work by studying other algebraic structures.

References

- [1] D. Molodtsov, Soft set theory-First results, Computers Mathematics Applications, 37(4-5)(1999), 19-31.
- [2] P.K. Maji, A.R. Roy, R. Biswas, An application of soft sets in a decision making problem, Computers Mathematics Applications, 44(2002), 1077-1083.
- [3] D. Chen, E.C.C. Tsang, D.S. Yeung, X. Wang, The parameterization reduction of soft sets and its applications, Computer Mathematics Applications, 49(2005), 757-763.
- [4] P.K. Maji, R. Biswas, A.R. Roy, Soft set theory, Computers Mathematics Applications, 45(2003), 555-562.

- [5] Z. Kong, L. Gao, L. Wang and S. Li, The normal Parameter Reduction of Soft Sets and Its Algorithm, *Computers and Mathematics with Applications*, 56 (1) 3029-3037.
- [6] J. E. Macias Diaz, On modules which are isomorphic to relatively divisible or pure submodules of each other, *Quaestiones Mathematicae*, 38 (6), 2015.
- [7] J. Abuhlail, C. Lomp, Topological lattices and their applications to modules, *Journal of Algebra Its Applications*, 15 (3), (2015).
- [8] R. Ameri, R. Sadeghi, Gamma modules, *Ratio Mathematica* 20, 127–147 (2010).
- [9] A. H. Rezaei, B. Davvaz, Tensor product of gamma modules, *Article in Afrika Mathematica*, 26 (7), 1601- 1608, (2015).
- [10] Q-M. Sun, Z-L. Zhang and J.Liu, Soft sets and modules (In Guoyin Wang, Tian-rui Li, Jerzy W. Grzymala-Busse, Duoqian Miao, Andrzej Skowron, Yiyu 54 Yao Eds.) *Rough sets and knowledge technology, RSKT, Proceedings, Springer*, 403-409, (2008).
- [11] D.Xiang, Soft module theory, *IEEE*, 2013, 10th International Conference on Fuzzy Systems and Knowledge Discovery.
- [12] T.Shah and S.Medhit, Primary decomposition in a soft ring and soft module, *Iranian Journal of Science & Technology*, 38A3 (Special issue-Mathematics) 311-320, (2014).
- [13] A.Erami, A.Hasankhani and A.B.Saeid, MV-modules in view of soft set theory, *Çankaya University Journal of Science and Engineering*, Volume 13, No. 1 (2016) 001–015.
- [14] M. I. Ali, F. Feng, X. Liu, W.K. Min, and M. Shabir, On some new operations in soft set theory, *Comput. Math. Appl.*, 57 (9), 1547-1553, (2009).
- [15] D. Pei and D. Miao, From soft sets to information systems, In *proceedings of Granular Computing* (Eds: X. Hu., Q. Liu, A. Skowron, T.Y.Lin, R.R. Yager, B. Zhang) *IEEE*, 2, 617-621.
- [16] Z. Xiao, Y. Li, B. Zhong and X.Yang, Research on synthetically evaluating method for business competitive capacity based on soft set, *Statical Research*, 52-54.
- [17] H. Aktas, N. Çağman, Soft sets and soft groups, *Inform. Sci.* 177 (2007), 2726- 2735.
- [18] U. Acar, F. Koyuncu, B. Tanay, Soft sets and soft rings, *Computers and Mathematics with Applications*, 59 (11), 2010, 3458- 3463.